

On the gamma-ray spectra radiated by protons accelerated in SNR shocks near molecular clouds: The case of SNR RX J1713.7-3946

M.A. Malkov¹, P.H. Diamond¹ and R.Z. Sagdeev²

Abstract

Cosmic rays (CRs) are thought to be accelerated in SNRs. The most favorable situation for proving that the main, hadronic CR component is accelerated there is when CRs interact with dense gases, such as molecular clouds (MC) which surround the SN shock. Here, a new mechanism of spectrum formation in partially ionized gases near SNRs is proposed. Using an analytic model of nonlinear diffusive shock acceleration, we calculate the spectra of protons and estimate the resulting γ -ray emission occurring when the SNR shock approaches a MC. We show that the spectrum develops a break in the TeV range and that its GeV component is suppressed. These modifications to the standard theory occur because of the proximity of the partially ionized MC-gas and because of the physics of particle and Alfvén wave propagation inside the gas. Possible applications of the new spectra to the recent CANGAROO and HESS observations of the SNR RX J1713.7-3946 are discussed.

Subject headings: acceleration of particles—cosmic rays —shock waves—supernova remnants—turbulence

1. Introduction

Cosmic rays (CRs) travel to us through the chaotic magnetic field of the Galaxy. Therefore, the only way to connect them to their accelerator is through the on-site radiation. Accelerated electrons have already been detected in supernova remnant (SNR) shocks (Koyama et al. 1995). However, electrons comprise only 1-2% of the CR intensity above 4-5 GeV. Therefore, distinguishing between leptonic and hadronic origin of the observed TeV emission from SNRs is a key to the proof of the supernova origin of CRs. Simultaneous monitoring of X- and γ - energy bands is an indispensable tool for that purpose. The same TeV *electrons* radiate in X- and γ -rays via synchrotron and inverse Compton (IC) mechanisms, respectively (*e.g.*, Sturmer et al. 1997; Reynolds

¹University of California at San Diego, La Jolla, California 92093-0319, USA; E-mail: mmalkov@ucsd.edu

²University of Maryland, College Park, Maryland 20742-3280, USA

1998). Accelerated *protons*, can be detected *only* near their own (most interestingly TeV) energy band through their interaction with an ambient gas. Therefore, the “smoking gun” for *proton acceleration* should be the γ -ray emission, without an X-ray emission that can be identified as the electron synchrotron radiation. Molecular clouds (MC) adjacent to the SNR will dramatically enhance proton visibility (*e.g.*, Aharonian et al. 1994; Drury et al 1994).

However, to conclusively detect acceleration of super-TeV nuclei turned out to be a very difficult task. Despite extensive search campaigns (*e.g.*, Buckley et al. 1998; Völk 2000), only three SNRs have shown detectable TeV emission so far (Tanimori et al. 1998; Muraishi et al. 2000; Aharonian et al. 2001; Enomoto et al. 2002). Nevertheless, a signature of protons accelerated to super-TeV energies was reportedly discovered in the SNR RX J1713.7-3946 by the CANGAROO team (Enomoto et al. 2002). Recently, this remnant has been confirmed as a TeV source, with significantly reduced systematic and statistical errors in the range 1-10 TeV, by the HESS collaboration (Aharonian et al. 2004). This remnant is of a shell type, typical for the major acceleration models, and rather abundant in the Galaxy. Therefore, after some verification of the data and analysis, it could serve as a direct evidence of the CR - SNR connection.

A number of groups (*e.g.*, Butt et al. 2002; Reimer and Pohl 2002) reanalyzed the data of Enomoto et al. (2002) and claimed that the nucleonic interpretation is “highly unlikely” because of an alleged inconsistency with the commonly assumed first order Fermi acceleration theory. We argue below that if (as it is assumed for this SNR by Butt et al. 2001; Enomoto et al. 2002), some of the accelerated particles begin to interact with a partially ionized dense gas such as a MC in the *northwestern* rim of the remnant, the standard acceleration model is not applicable. In particular, the nonlinearity of the acceleration process as well as Alfvén wave evanescence in the MC are essential. Therefore, the analyses by both sides of the controversy are oversimplified and we believe that a conclusive case for or against nucleonic origin of the emission has not been made. Moreover, a recent analysis by Pannuti et al. (2003) indicates that the standard acceleration model applied to TeV electrons does not adequately fit the TeV data, either, unless the X-ray emitting filaments have an unreasonably low filling factor of 10^{-3} . Lazendic et al. (2004) have increased it to 10^{-2} by allowing a smoother spectrum cut-off.

These analyses indicate that modifications to the accelerated particle spectra are required. Namely, a low-energy cut-off above 100 GeV and/or a *spectral break* would support the nucleonic scenario (Reimer and Pohl 2002) while models with an exponential cut-off rather than a break provide only a bad fit. Such modifications are suggested below.

The controversy about the TeV observations of the SNR RX J1713.7-3946 is fundamental and will clearly impair understanding of future observations of SNRs nearby MCs. Therefore, in this Letter we revisit shock acceleration theory and include the following physical phenomena in the model: (i) the nonlinearity of the acceleration process, *i.e.*, the modification of the flow by

accelerated particles (ii) a position dependent *low-energy cut-off* of accelerated particles ahead of the shock, which is *well known* to be present in analytic solutions for the shock acceleration problem. Note that this cut-off is usually ignored in shock acceleration models since only the downstream solution is considered. When the dense gas is present upstream of the shock this cut-off should be included in the calculation of γ -radiation. (iii) impairment of CR confinement in a dense gas (MC) due to non-propagation of some Alfvén waves. The latter phenomenon produces a break (curvature) in the particle and radiation spectra in the TeV energy range and, in combination with (i), results in a spectral slope significantly different from the usual predictions of linear theory supplemented with the high energy cut-off, which was used in the analyses of Enomoto et al. (2002) and Reimer and Pohl (2002), for example.

2. Low-energy cut-off and the absence of thermal x-ray emission

The diffusive shock acceleration (DSA) mechanism operates by scattering charged particles off magnetic irregularities (Alfvén waves) across a shock, such as a SN blast wave (*e.g.*, Blandford and Eichler 1987). In the linear acceleration regime, the momentum distribution is a power-law, $p^2 f_0(p) = Cp^{-s}$ where C is a normalization constant and the index $s = (r + 2)/(r - 1)$ depends only on the shock compression ratio r . As soon as the flow is disturbed by the pressure of accelerated particles, the spectrum becomes more complex. It flattens toward higher energies and may become steeper at lower energies. The spectrum $f_0(p)$ is the *downstream* spectrum, which is coordinate independent in both the linear and nonlinear regimes. Here the upstream spectrum is more important since the interaction with the adjacent MCs starts *upstream* where a low energy *cut-off* occurs. Indeed, the solution upstream, valid for both the linear and nonlinear regimes, reads (*e.g.*, Malkov and Drury 2001)

$$f(p, x) = f_0(p) \exp \left[-\frac{\alpha}{\kappa} \int_0^x u dx \right]. \quad (1)$$

Here $f_0(p)$ is the spectrum downstream, $\alpha(p) \sim 1$. The x coordinate points upstream ($x > 0$) from the shock ($x = 0$) and $u(x) > 0$ is the speed of the plasma flowing into the shock ($u(x) \sim V_{shock}$). Now, $\kappa(p)$ grows with p , most likely linearly, $\kappa \approx \kappa' p$. According to eq.(1), there exists a CR precursor, which is of the length $l_{CR} = \kappa(p_{max})/V_{shock}$. Furthermore, high energy particles diffuse ahead of low energy particles so the spectrum (1) has an exponential *low-energy cut-off* at

$$p_{min}(x) \approx \frac{1}{\kappa'} \int_0^x u dx \sim V_{shock} x / \kappa'. \quad (2)$$

For the Bohm diffusion we have

$$p_{min}/mc \approx (V_{shock}/c) B_{\mu} (x/10^{12} cm) \quad (3)$$

where B_{μ} is the magnetic field in μG . This low-energy decay can explain why the same MC can be visible in the TeV energy range and invisible in the GeV range. Let the near-most edge of an adjacent MC be upstream of the shock at a distance $x = x_{MC}$, such that $1 GeV/c < p_{min}(x_{MC}) < p_{max}$. This low-energy cut-off p_{min} is shown in Fig.1¹ that will be discussed below. The last condition means that the leading edge of the CR precursor (filled with TeV protons) has already penetrated into the MC, but the sub-shock itself (with the GeV particles ahead of it) still has not. Obviously, the CANGAROO and HESS could detect the proton TeV emission while the EGRET could not, because of the low density inside the wind bubble the sub-shock is located in. The strong X-ray emission expected when the sub-shock crashes into the dense MC will not yet be visible, either.

3. High-energy spectral break

In the presence of weakly ionized dense gas the particle and emission spectra undergo significant modifications in the TeV energy range, where a spectral break can form. According to Eq. (1) the accelerated particles occupy an extended precursor of the size

$$l_{CR} \sim \kappa(p_{max})/V_{shock} > r_g(p_{max})c/V_{shock} \quad (4)$$

ahead of the shock. Here r_g is the particle gyroradius. One can estimate l_{CR} as $l_{CR} \sim (p_{max}/mc) (c/V_{shock}) B_{\mu}^{-1} 10^{12} cm$. Thus, the shock precursor may be as long as $10^{18} cm$ for $p_{max} \sim 10 TeV$. Therefore, the accelerated particles start to interact with a MC before it becomes significantly ionized by the shock wave or the ionizing precursor (Drane and McKee 1993). Hence, they propagate in the MC under conditions of strong ion-neutral collisional damping of the self-generated Alfvén waves that are needed to confine accelerated particles. More importantly, there is a gap in wavenumber space at $k_1 < k < k_2$ where the waves do not propagate (Kulsrud and Pearce 1969; Zweibel and Shull 1982). Here, it is critical to realize that waves *literally do not exist*, as opposed to simply being damped by collisions. The above wave evanescence range is bounded by $k_1 = v_{in}/2V_A$ and $k_2 = 2\sqrt{\rho_i/\rho_0}v_{in}/V_A$, where $\rho_i/\rho_0 \ll 1$ is the ratio of the ion to neutral mass density, v_{in} is the ion-neutral collision frequency and $V_A = B/\sqrt{4\pi\rho_i}$ is the Alfvén speed. The resonance condition for the wave generation and particle scattering off them is $kp_{\parallel}/m = \pm\omega_c$,

¹We have chosen the following values for the parameters in eq.(3): $V_{shock} = 1000 km/sec$, $B_{\mu} = 3$, and the distance from the sub-shock to the MC, $x = x_{MC} \approx 7 \cdot 10^{15} cm$. Note that for the visibility of the TeV and higher energy protons, it is sufficient that $x_{MC} < 10^{17} cm$.

where $\omega_c = eB/mc$ and $p_{\parallel}$ is the parallel (to the magnetic field) component of particle momentum. Therefore, particles having $p_1 < |p_{\parallel}| < p_2$, where

$$p_1 = 2V_A m \omega_c / v_{in}, \quad p_2 = \frac{1}{4} \sqrt{\frac{\rho_0}{\rho_i}} p_1 > p_1, \quad (5)$$

are not scattered by any waves, so that the confinement of all particles with $|p_{\parallel}| > p_1$ is dramatically degraded. We assume that the gap in $p_1 < p_{\parallel} < p_2$ is sufficiently broad, so that resonance broadening (*e.g.*, Achterberg 1981) does not bridge it. We also assume that there is no significant background turbulence at the scales $\lambda > 2\pi/k_1$, that could reduce particle diffusivity to a limit which is not significantly higher than the Bohm value. For the parameters used in our calculations below, $k_1 \sim 10^{-14} \text{cm}^{-2}$. Then, particles with $|p_{\parallel}| > p_1$ will escape from the MC upstream of the shock along the magnetic field at a speed comparable to c or, at least, their confinement time to the shock will be much shorter than those with $|p_{\parallel}| < p_1$. Only the latter particles will generate π^0 mesons and γ emission efficiently in the MC. Their pitch angle distribution for $p > p_1$ is, however, limited to the interval $|\mu| < p_1/p$. Therefore, the contribution to the γ emission of particles with momentum $p > p_1$ is reduced by the phase volume filling factor $\mu_{crit} = p_1/p$. The resulting emission spectrum is thus one power steeper as compared to the standard calculations (*e.g.*, Berezhinsky and Ptuskin 1989; Drury et al 1994; Naito and Takahara 1994) based on the isotropic particle distribution. The latter yields the γ spectrum which reproduces that of the particles up to about $0.1cp_{max}$ where it declines. Hence, the energy spectrum of γ emission must be the same as that of the particles, $\propto \epsilon_{\gamma}^{-s}$, for the photon energy $\epsilon_{\gamma} < \epsilon_{br} \sim 0.1cp_1$ (particle momenta $p < p_1$), and it must scale as $\propto \epsilon_{\gamma}^{-s-1}$ for $\epsilon_{\gamma} \gtrsim \epsilon_{br}$ up to about $0.1cp_{max}$ (particle momenta $p_1 < p < p_{max}$). The break energy ϵ_{br} can be estimated using Eq. (5) for p_1 by substituting $v_{in} = n_0 \langle \sigma V \rangle$, where σ is the cross-section of the ion-neutral collisions and V is the collision velocity averaged over a thermal distribution. Using an approximation of Kurlsrud and Cesarsky (1971) for $\langle \sigma V \rangle$, p_1 can be estimated as

$$p_1/mc \simeq 16B_{\mu}^2 T_4^{-0.4} n_0^{-1} n_i^{-1/2}. \quad (6)$$

where T_4 is measured in the units of $10^4 K$, n_0 and n_i (number densities corresponding to the mass densities ρ_0 and ρ_i) -in cm^{-3} .

To illustrate these results we consider three different acceleration regimes, corresponding to the three substantially different shock compression ratios r , shown by the points 1-3 in the inset to Fig.1. (Here r means a total shock compression, that is adiabatic compression across the precursor times the subshock jump.) As it can be seen from this plot, the shock compression is difficult to calculate accurately due to its sharp dependence on the injection rate (usually not well known) and other parameters, such as the maximum momentum. We discuss this spectrum in more detail in the next section.

4. Discussion and Conclusions

In this paper we have considered the *upstream* particle spectrum since this is necessary for interpretation of cases where a shock is expanding into a low-density pre-supernova wind bubble and is approaching a denser material such as a swept-up shell or a MC. Perhaps the most discussed SNR of this kind is the RX J1713.7-3946 claimed by the CANGAROO team to be a long anticipated proton super-TeV accelerator (Enomoto et al. 2002). Significant part of the TeV emission detected by the CANGAROO and HESS (Aharonian et al. 2004) instruments comes from the northwestern rim of the remnant where the interaction with a molecular cloud is believed to take place (*e.g.*, Slane et al. 1999; Hiraga et al. 2004). Therefore these observations are pertinent to the subject of this letter.

To demonstrate that the observed TeV spectra are consistent with the mechanisms suggested in this paper, we put the calculated upstream *proton* spectra on one plot with the CANGAROO/HESS and EGRET *photon* fluxes that would be radiated by these protons. Since the emission generated by protons has typically an order of magnitude higher energy, we up-shifted the *observed* emission energy accordingly in Fig. 1. As the spectrum between $\text{GeV}/c < p < p_1$ is considerably flatter than in the case of the linear theory, its magnitude at $p = p_1$ is an order of magnitude higher for the same injection rate. This lowers the ambient gas density required to produce the same TeV emission as computed by Enomoto et al. (2002) using the linear spectrum. In addition, this allows to accommodate the EGRET points, should they belong to the same object (Reimer and Pohl 2002). However, because of the controversy about the distance to the remnant (6 vs 1 kpc) (Slane et al. 1999; Fukui et al. 2003) and other uncertainties, it is impossible to present detailed normalized fits of the photon flux in this short letter.

The form of the spectrum is marked by increasing the spectral index by one at the break momentum $p = p_1$, Fig. 1. This is crucial to fitting the observed spectra without imposing a specific form and value of the energy cut-off. The latter procedure would give only a poor fit (*e.g.*, Reimer and Pohl 2002). Note that there are two CANGAROO points that are not in a good agreement with both the HESS and with the theoretical spectrum. It should be noted, however, that any disagreement between the three sets of spectral points is substantially enhanced in this particular spectrum format (the particle phase density is multiplied by p^2). Except for these two points the agreement is remarkably good. Interestingly enough the agreement is excellent for a subset of HESS points with the lowest and highest energy points excluded.

Unlike the power-law index, it is more difficult to constrain the parameters determining the position of the break on the spectrum given by Eq. (6) due to the poor information about the target gas for $p - p$ reactions (MC). For the particular case shown in Fig.1, we have chosen such combination of parameters in Eq. (6) that p_1 amounts to $p_1 \sim 1.8\text{TeV}/c$. MCs in general are known to be “clumpy” with the inter-clump gas density of $5\text{-}25\text{ cm}^{-3}$ and a less than 10% filling

factor (see, *e.g.*, Chevalier 1999; Bykov et al. 2000, and references therein) . The results shown in Fig. 1 are obtained for the following values of parameters in eq.(6): $B_\mu \approx 14$, $T_4 \approx 0.01$, $n_0 \approx 23$, and $n_i \approx 0.23$ which yields $p_1/mc \approx 1.8 \cdot 10^3$. By contrast, in the model of Ellison et al. (2001) much lower target density is assumed since its authors do not consider the possibility that the CR precursor reach the dense gas while the shock itself is still in the wind bubble. Therefore they concluded that the $p - p$ reactions do not contribute significantly to the TeV emission because of the lack of target protons. In contrast to the conclusion of Ellison et al. (2001) and to that of Enomoto et al. (2002) the most recent HESS observations allow their authors (Aharonian et al. 2004) to assume that both protons and electrons contribute to the TeV emission significantly.

Note that a complementary mechanism of suppression of the low-energy (GeV) emission has been suggested earlier by Aharonian and Atoyan (1996) and discussed recently in the context of SNR RX J1713.7-3946 by Uchiyama et al (2003). This mechanism requires an impulsive release of accelerated particles at some distance from a MC (target) and subsequent (energy dependent) diffusive propagation of CRs to the target. By contrast, the mechanism discussed in this paper is based on a quasi-stationary solution of the acceleration problem (Eq. [1]) which implies that the accelerated particles are bound to the propagating shock front via self-generated Alfvén waves. It is clear, however, that in both cases the suppression of the low-energy emission is based on a slower diffusion of low-energy particles. Note that Lazendic et al. (2004) also discussed qualitatively the same effect. It is also important to mention that GeV emission may not be suppressed all over the remnant, and was demonstrated by Butt et al. (2001) to be likely of proton origin. Obviously, the spectral modifications considered in this letter are equally applicable to electrons of similar rigidity.

It should be pointed out that our model in its current version does not provide a complete fit to the data, particularly in the X- and radio energy bands. The TeV-GeV fluxes are given in arbitrary units, so only the *form* of the spectrum (including possible GeV signal) is provided here. A complete fit would require the convolution of the proton spectrum with appropriate emissivities given the densities and the distance. The thermal peak in Fig.1 is a Maxwellian, approximately normalized relative to the high-energy part of the spectrum, in accordance with the calculated downstream temperature without turbulent heating. More accurate calculation of the thermal peak would require adequate injection model and turbulent gas heating in the precursor (*e.g.*, Berezhko et al. 1996; Malkov and Drury 2001; Kang et al. 2002). Such calculations of the thermal X-ray emission should be tested specifically against very low observed upper limits. One *potentially* observable prediction of our model is the break in the pitch angle averaged particle spectrum at $p = p_1$ that would be produced by almost instant losses along the field lines of all particles with $|p_{||}| > p_1$ once they enter a partially ionized gas with significant ion-neutral collisions. This break may very well be in the currently “obscure” energy range between the EGRET and the ground based Cherenkov telescope energy bands. The next generation of space gamma telescopes, such

as GLAST, will be able to explore this energy range.

The principal results of this letter, which are the spectrum softening by one degree above the spectral break, and the possibility of a low energy cut-off clearly show that these physical phenomena need to be included in the models to conclusively differentiate between the nucleonic and leptonic sources of the TeV emission from the remnants nearby molecular clouds such as the RX J1713.7-3946. Previous claims to the contrary have ignored necessary refinements in the DSA theory beyond the level of test particle theory.

We thank Felix Aharonian and Heinz Völk for helpful discussions as well as an anonymous referee provided useful comments on the manuscript. Support under NASA grant ATP03-0059-0034 and under U.S. DOE grant No. FG03-88ER53275 is gratefully acknowledged.

REFERENCES

- Achterberg, A. 1981, A&A, **98**, 161
- Aharonian, F.A., Drury, L.O’C. and Völk, H.J. 1994, A&A, 285, 645
- Aharonian, F.A., Atoyan, A.M. 1996, A&A, 309, 917
- Aharonian, F. A. *et al.* 2001, A&A, 370, 112
- Aharonian, F. A. *et al.* 2004, Nature, 432, 75
- Berezinsky, V.S. & Ptuskin, V.S. 1989 ApJ, 340, 340
- Berezhko, E. G., Yelshin, V., & Ksenofontov, L. 1996, Sov. Phys. JETP , 82, 1
- Blandford, R.D. & Eichler, D., 1987, Phys. Rep., 154, 1
- Buckley, J. H. *et al.* , 1998, A&A, 329, 639
- Butt, Y.M. *et al.* 2002, Nature, 418, 499
- Butt, Y.M. *et al.* 2001, ApJ, 562, L167
- Bykov, A.M. *et al.* , 2000, ApJ, 538, 203
- Chevalier, R.A., 1999, A&A, 511, 798
- Draine, B.T., McKee C.F., 1993, Annu. Rev. Astron. Astrophys., 31, 373

- Drury, L. O’C, Aharonian, F. A. & Völk, H. J., 1994, A&A, 287, 959
- Drury, L. O’C, Duffy, P. Kirk, J. K., 1996, A&A, 309, 1002
- Ellison, D.C., Slane, P. & Gaensler, B.M. 2001, ApJ, 563, 191
- Enomoto, R. *et al.* 2002, Nature, 416, 823
- Fukui, Y. *et al.* 2003, PASJ, 55, L61
- Hartman, R.C. *et al.* 1999, ApJS, 123, 79
- Hiraga, J.S. *et al.* , 2003, astro-ph/0407401
- Kang, H., Jones, T.W. & Gieseler, U.D.J. 2002, ApJ579, 337
- Koyama, K. *et al.* 1995, Nature, 378, 255
- Kulsrud, R. & Pearce, W.P. 1969, ApJ, 156, 445
- Kulsrud, R.M. & Cesarsky, C.J. 1971, ApJ, 8, 189
- Lazendic, J.S. *et al.* , 2004, ApJ, 602, 271
- Malkov, M. A. & Drury L.O’C., 2001, *Rep. Progr. Phys.* , 64, 429
- Muraishi, H. *et al.* 2000, A&A, 354, L57
- Naito, T. & Takahara, F. 1994, J. Phys. G20, 477
- Pannuti, T.G., Allen, G.E., Houck, J.C. and Sturmer, S.J. 2003, ApJ, 593, 377
- Reynolds, S.P. 1998 ApJ, 493, 375
- Reimer, O. & Pohl, M. 2002, A&A, 390, L43
- Slane, P. *et al.* 1999, ApJ525, 357
- Sturmer, S.J., Skibo, J.G., Dermer, C.D. & Mattox J.R. 1997, ApJ490, 619
- Tanimori, T. *et al.* 1998, ApJ, 497, L25
- Uchiyama, Y., Aharonian, F.A., and Takahashi, T., 2003, A&A, 400, 567
- Völk, H.J. 2000, in AIP Conference Proceedings, 515, 197
- Zweibel, E., and Shull, J.M. 1982, ApJ, 259, 859




Fig. 1.— Inset: response of the shock structure to the particle acceleration. The flow compression ratio r is shown as a function of particle density n_{CR} injected into the acceleration from the thermal distribution calculated for two maximum momenta $p_{\max}$ and the shock Mach number $M=80$. The injection parameter $v \sim (cp_{\text{inj}}/mV_{\text{shock}}^2)n_{CR}/n_1$, where n_1 is the plasma density upstream of the shock. The details of these calculations and those of the spectra shown in the main figure can be found in (Malkov and Drury 2001). Main figure: spectra of accelerated particles behind the shock front, for $p_{\max}/mc = 10^5$. Three solutions correspond to the points 1-3 on the compression-injection diagram in the inset. Note that the test particle (linear) spectrum would be a horizontal line. The break energy (momentum) is at $p/mc = 1.8 \cdot 10^3$. The CANGAROO-II (Enomoto et al. 2002) and HESS points are adopted from Aharonian et al. (2004) while EGRET (Hartman et al. 1999) points— from Reimer and Pohl (2002). Vertical lines indicate: the injection momentum (separating thermal and nonthermal particles), the 1.2 GeV proton energy threshold of π^0 production, the possible low-energy cut-off $p_{\text{min}}(x)$ and the spectral break p_1 (see text).